

A user's guide for *atompaw* code

Marc Torrent

Commissariat à l'énergie atomique. BP12. 60680 Bruyères-le-Chatel – France

Natalie A. W. Holzwarth

Wake Forest University. Winston-Salem, NC 27109 - USA

Revised august 27th, 2007

Compatible with atompaw v2.2 and later

METHOD FOR PAW DATASET GENERATION	2
HOW TO USE <i>ATOMPAW</i>	3
INPUT FILE FOR <i>ATOMPAW</i>	4
DETAILED DESCRIPTION OF KEYWORDS	5
1. Atomic all-electron computation	5
2. Partial-wave basis generation	7
3. Test configurations	10
EXAMPLES	11
ADVICE FOR USE	13
Short write-up	13
Detailed write-up	14

Source code URLs : <http://pwpaw.wfu.edu>, <http://www.abinit.org/PAW/>

Contact emails : marc.torrent@cea.fr, natalie@wfu.edu

Method for PAW dataset generation

PAW calculations require, for each atomic species, a set of basis (partial-waves) and projectors functions plus some additional atomic data stored in a *PAW dataset*. A PAW dataset has to be generated in order to reproduce atomic behavior as accurate as possible while requiring minimal CPU and memory resources in executing the DFT code for the crystal simulations. These two constraints are conflicting.

The PAW dataset generation is done according the following procedure:

All parameters that should be given in an *atompaw* input file are in **bold**.

- 1- Choose and define the concerned chemical species: **name** and **atomic number**.
- 2- Solve the atomic all-electrons problem in a given atomic configuration. The atomic problem is solved within the DFT formalism, using an **exchange-correlation functional** and either a Schrödinger (default) or **scalar-relativistic approximation**. It is a spherical problem and it is solved on a **radial grid**. Other approximations can be given (as, for example, the behavior of the **nuclear potential**). The atomic problem is solved for a given **electronic configuration** that can be an ionized/excited one.
- 3- Choose a set of electrons that will be considered as frozen around the nucleus (**core electrons**). The others electrons are valence ones and will be used in the PAW basis. The **core density** is then deduced from the core electrons wave functions. A **smooth core density** equal to the core density outside a given r_{core} **matching radius** is computed.
- 4- Choose the **size of the PAW basis** (number of partial-waves and projectors). Then choose the partial-waves included in the basis. The later can be **atomic eigen-functions** related to valence electrons (bound states) — in fact this is mandatory with *atompaw* — and/or **additional atomic functions**, solution of the wave equation for a given l **quantum number** at arbitrary **reference energies** (unbound states).
- 5- Generate **pseudo partial-waves** (smooth partial-waves build with a **pseudization scheme** and equal to partial-waves outside a given r_c **matching radius**) and associated **projector functions**. Pseudo partial-waves are solutions of the PAW Hamiltonian deduced from the atomic Hamiltonian by pseudizing the effective potential (a local **pseudopotential** is built and equal to effective potential outside a r_{yloc} **matching radius**). Projectors and partial-waves are then orthogonalized with a chosen **orthogonalization scheme**.
- 6- Build a *compensation charge density* used later in order to retrieve the total charge of the atom. This compensation charge density is located inside the PAW spheres and based on an analytical **shape function** (which analytic form and **localization radius** r_{shape} can be chosen).
- 7- Eventually, if desired, test the resulting PAW dataset on several **electronic test configurations**.

How to use *atompaw*

- 1- Compile *atompaw*:
Edit [make.inc](#) in the *atompaw* directory according to local specifications.
Type : `make`
- 2- Edit an input file in a text editor (content of input is explained in the following).
- 3- Run *atompaw*: `atompaw < inputfile`

Partial-waves, PS partial-waves and projectors are given in [wfn.i](#) files.

Logarithmic derivatives from atomic Hamiltonian and PAW Hamiltonian resolutions are given in [logderiv.l](#) files.

A summary of the atomic all-electrons computation and PAW dataset properties can be found in the [Atom_name](#) file (*Atom_name* is the first parameter of the input file).

Resulting PAW dataset is contained in:

- [Atom_name.atomicdata](#) file
Specific format for *pw paw* code
- [Atom_name.XCfunc.xml](#) file
Normalized *xml* file according to specifications from <http://dcwww.camp.dtu.dk/campos/pawxml.pawxml.xhtml>.

A PAW dataset file in a format suitable for *abinit* code can be obtained using *Atompaw2Abinit* tool (<http://www.abinit.org/PAW/AtomPAW2Abinit-Manual-html/HolzwarthPAW.htm>).

Additional details can be found in:

- “Notes for revised form of *atompaw* code”
<http://www.wfu.edu/~natalie/papers/pwpaw/notes/atompaw/atompawEqns.pdf>
- “Part I manuscript.pdf”
[http://dx.doi.org/10.1016/S0010-4655\(00\)00244-7](http://dx.doi.org/10.1016/S0010-4655(00)00244-7)
(be careful: some obsolete chapters inside).

Input file for *atompaw*

In *red*, *mandatory* arguments
In *green*, *optional* arguments

Keywords are in normal font
Numbers are in italics

<pre> Atom_name Z XC_functional rel_keyword nucleus_keyword loggrid_keyword gridsize <i>n_s^{max}</i> <i>n_p^{max}</i> <i>n_d^{max}</i> <i>n_f^{max}</i> <i>n_g^{max}</i> n l occ_{nl} n l occ_{nl} ... 0 0 0 c or v c or v c or v c or v c or v ... </pre> <p>One line for each empty or partially occupied (n,l) state</p> <p>One line for each (n,l) state l=0 states first then l=1 states...</p>	<p>Atomic all-electrons computation</p>
<pre> <i>l_{max}</i> <i>r_{paw}</i> <i>r_{shape}</i> <i>r_{vloc}</i> <i>r_{core}</i> Y E_{ref} n Y E_{ref} n ... Y E_{ref} n projector_keyword ps_scheme ortho_scheme shapefunction tol <i>l_{loc}</i> <i>E_{loc}</i> <i>Vloc_scheme</i> rc₁ rc₂ ... rc_{basis_size} </pre> <p>Repeated for each additional l=0 partial-wave</p> <p>Repeated for each additional l=1 partial-wave</p> <p>Repeated for each additional l=l_{max} partial-wave</p> <p>One paragraph for each 0 ≤ l ≤ l_{max}</p> <p>If Projector_keyword ≠ "Bloechl"</p> <p>One line for each partial-wave</p>	<p>Partial-waves basis generation</p>
<pre> 1 n l occ_{nl} n l occ_{nl} ... 0 0 0 0 </pre> <p>One line for each empty or partially occupied (n,l) state</p> <p>As many times as desired</p>	<p>Tests configurations</p>

Detailed description of keywords

1. Atomic all-electron computation

Atom_name Z

Atom_name:
Symbol of atomic specie

Z:
Atomic number (total number of electrons)

XC_functional rel_keyword nucleus_keyword grid_keyword gridsize r_{max} r_{match}

XC_functional:
Name of exchange-correlation functional used in DFT atomic configuration resolution

Possible values:

- LDA-PW** for Perdew-Wang (92) LDA functional [PRB 45, 13244 (1992)]
- GGA-PBE** for Perdew-Burke-Ernzerhof (96) GGA functional [PRL 77, 3865 (1996)]

rel_keyword:
Relativistic approximation used to solve atomic wave equation

Possible values:

- nonrelativistic**: solve non-relativistic Schrödinger equation.
- scalarrelativistic**: solve scalar-relativistic wave equation (Koelling-Harmon-like equation). [J. Phys C 10, 3107 (1977)]

Default (if missing) is **nonrelativistic**

nucleus_keyword:
Option governing the form of the potential near $r \rightarrow 0$ during atomic wave equation resolution

Possible values:

- point-nucleus**: solve atomic wave equation, assuming point potential for $r \rightarrow 0$ ($V(r) = -2Z/r$).
- finite-nucleus**: solve atomic wave equation, assuming finite nucleus potential for $r \rightarrow 0$ ($V(r) = -2Z \operatorname{erf}(r/RR)/r$), where RR is a nuclear size parameter.

Default (if missing) is **point-nucleus**

grid_keyword gridsize r_{max} r_{match} :
Options governing the analytic form of the radial grid used in atompaw

Analytical form of the grid is determined by grid_keyword
Its step and size can be defined by gridsize, r_{max} and r_{match} .

Possible values for grid_keyword:

- lineargrid**: use a linear grid : $r_i = h(i-1)$
- loggrid**: use a logarithmic grid : $r_i = (h/Z) \cdot (\exp[h(i-1)] - 1)$

Default (if missing) is **lineargrid**

Additional (optional) arguments:

gridsize:
Define the number of points in the grid.
Default (if missing) is **20001** when grid is linear
2001 when grid is logarithmic

r_{max} :
Optional argument, but if present must follow gridsize in the input.
Define the maximum radius of the grid.
Default (if missing) is **50. a.u.** when grid is linear
80 a.u. when grid is logarithmic

r_{match} :
Optional argument, but if present must follow r_{max} in the input.
This changes the usage of gridsize so that the value of r_{match} defines an explicit grid point by adjusting the step size (h) so that there are gridsize grid points between 0 and r_{match} .
A typical value for r_{match} is the PAW radius, r_{paw} (in order to keep it constant when the grid size changes).
The grid is then continued to the first point $r_i \geq r_{max}$.
Default (if missing) is r_{max}

n_s^{\max} n_p^{\max} n_d^{\max} n_f^{\max} n_g^{\max}

n_l^{\max} :
 Maximum n quantum number for l electrons
 Example:
 For Nickel ($1s^2 2s^2 2p^6 3s^2 3p^6 3d^8 4s^2$), enter:
 4 3 3 0 0

n l occ_{nl}
 n l occ_{nl}
 ...
 0 0 0

One line for each empty or partially occupied (n,l) state

For each electronic shell of the atomic species, enter a line with:
 n,l : quantum numbers of the shell
 occ_{nl} : electronic occupation of the shell
 Actually, only empty or partially occupied shells are needed; full shells can be omitted.
 Charged/excited configurations are (of course) accepted.
 A "0 0 0" (zero zero zero) line ends the configuration.
 Example:
 For excited Nickel ($1s^2 2s^2 2p^6 3s^2 3p^6 3d^{8.5} 4s^{1.5}$), simply enter:
 3 2 8.5
 4 0 1.5
 0 0 0



c or v
 c or v
 c or v
 c or v
 c or v
 ...

One line for each (n,l) state
 $l=0$ states first
 then $l=1$ states...

c or v :
 Core or valence characteristic of electronic shell
 For each electronic shell, enter a c or v keyword, which can be:
 c : the electronic shell is a CORE shell, frozen around the nucleus and included in the core density of the PAW data set.
 v : the electronic shell is a VALENCE shell containing valence electrons included in the PAW data set. In addition, note that the partial-wave associated with such a valence state will be NECESSARILY included in the PAW partial-waves basis (see below).
 Example:
 For Nickel ($1s^2 2s^2 2p^6 3s^2 3p^6 3d^8 4s^2$), with $3p^6 3d^8 4s^2$ in the valence, enter:

c	! 1s
c	! 2s
c	! 3s
v	! 4s valence
c	! 2p
v	! 3p valence
v	! 3d valence

2. Partial-wave basis generation



Definition of partial-waves basis elements:

By construction, the basis already contains each atomic wave function associated with a valence state (each wave function marked as "v" in the atomic all-electron configuration). These are "bound states".

To add additional basis elements ("unbound states"), proceed as follow:

For each l angular momentum (from 0 to l_{max}),

- 1- Enter y to add an additional partial-wave
- 2- Enter E_{ref} (real number, Rydberg units), reference energy used to build the partial-wave. The later is obtained by inverting the Schrödinger equation at energy E_{ref} and l angular momentum.

Go to point 1- to add another partial-wave associated with l

Or

Enter n

projector_keyword:

Option governing the scheme used to generate (smooth) PS partial-waves and associated projectors

Possible values:

Bloch [or **VNCT**]: use P. Blochl PS wave functions and projectors generation scheme [PRB **50**, 17953 (1994)]: A cutoff-function $k(r)=[\sin(\pi/r_{paw})/(\pi/r_{paw})]^2$ is used (in a Schrödinger-like equation) to deduce PS partial-waves. Projectors are then orthogonalized with a Gram-Schmidt procedure. In that case, ps_scheme and ortho_scheme keywords are ignored.

Vanderbilt [or **VNCTV**]: use a polynomial function to “pseudize” partial-waves and D. Vanderbilt projectors generation scheme [PRB **41**, 7892 (1990)]:The polynomial function used to “pseudize” partial-waves is identical as the one used when ps_scheme=**polynom** (see below) In that case, ps_scheme and ortho_scheme keywords are ignored.

custom: get PS wave functions according to ps_scheme keyword (see below) and projectors according to ortho_scheme keyword (see below).

ps_scheme:

Option governing the scheme used to generate (smooth) PS partial-waves when projector_keyword=**custom**

Possible values:

blochlps: use P. Blochl PS wave functions and projectors generation scheme [PRB **50**, 17953 (1994)]: a cutoff-function $k(r)=[\sin(\pi/r_c)/(\pi/r_c)]^2$ is used (in a Schrödinger-like equation) to deduce PS partial-waves. In that case ortho_scheme keyword has to be **gramschmidtortho**.

polynom: use a eighth degree polynomial function to “pseudize” partial-waves.

Below matching radius, PS wave function has the form: $\tilde{\varphi}_i(r) = r^{l+1} \cdot \sum_{m=0}^4 C_m r^{2m}$

polynom2 p q_{cut}: use a polynomial of degree $2p$ to “pseudize” partial-waves.

Below matching radius, PS wave function has the form: $\tilde{\varphi}_i(r) = r^{l+1} \cdot \sum_{m=0}^p C_m r^{2m}$

For $m \geq 4$, C_m coefficients are computed so that to minimize Fourier coefficients of PS partial-wave for $q > q_{cut}$ (*Fourier filtering*).

Defaults values of p and q_{cut} (if missing) are: $p=4$; $q_{cut}=10.0$

rrkj: use RRKJ scheme to get PS wave functions [PRB **41**, 1227 (1990)].

Below matching radius, PS wave function is a sum of 2 Bessel functions:

$$\tilde{\varphi}_i(r) = r \cdot (\alpha'_1 \cdot j_l(q'_1 r) + \alpha'_2 \cdot j_l(q'_2 r))$$

Default (if missing) is **bloch**lps

ortho_scheme:

Option governing the scheme used to generate and orthogonalize projectors when projector_keyword=**custom**

Possible values:

gramschmidtortho: use a Gram-Schmidt –like procedure to orthogonalize projectors and PS partial-waves.

vanderbiltortho: use D. Vanderbilt procedure to orthogonalize projectors and PS partial-waves (see [PRB **41**, 7892 (1990)]).

Default (if missing) is **gramschmidtortho**

Note that:

projector_keyword=**vanderbilt** is strictly equivalent to “**custom polynom vanderbiltortho**”

projector_keyword=**blochl** is equivalent to “**custom blochlps gramschmidtortho**”

AND all r_c (defined later) equal to r_{paw}

shapefunction tol:

Option governing the analytic form of shape functions $g_l(r)$ used in compensation density definition

Can be:

sinc: $g_l(r) = N \cdot r^l \cdot k(r)$ with $k(r) = [\sin(\pi r / r_{shape}) / (\pi r / r_{shape})]^2$
tol parameter is ignored and can be omitted

gaussian tol: $g_l(r) = N \cdot r^l \cdot k(r)$ with $k(r) = \exp(-(r/d)^2)$
d parameter is deduce so that $k(r_{shape}) = \mathit{tol}$
Default of *tol* parameter (if missing) is 10^{-4}

besselshape: $g_l(r) = \alpha_1^l \cdot j_l(q_1^l r) + \alpha_2^l \cdot j_l(q_2^l r)$ (see [PRB 59, 1758 (1999)])
tol parameter is ignored and can be omitted

Default (if missing) is **sinc**

l_{loc} E_{loc} **Vloc_scheme**

Vloc_scheme:

Option governing the scheme used to get $V^{PS}(r)$ (local) pseudopotential from all-electron effective potential $V_{eff}(r)$. Matching radius for pseudization is r_{vloc} .

Can be:

troulliermartins: use a norm-conserving Troullier-Martins scheme. A PS wave function is deduce from atomic one and chosen to have the form $\phi^{PS}(r) = r^{1+l_{loc}} \cdot \exp(p(r))$ for $r < r_{vloc}$ where p is an even 12th order polynomial. Then V^{PS} is deduced by inverting the wave equation at $l=l_{loc}$ and $E=E_{loc}$.

ultrasoft: use a pseudization scheme without norm conservation constraint. A PS wave function is deduce from atomic one and chosen to have the form $\phi^{PS}(r) = r^{1+l_{loc}} \cdot \sum_{m=0}^3 C_m r^{2m}$ for $r < r_{vloc}$. Then V^{PS} is deduced by inverting the wave equation at $l=l_{loc}$ and $E=E_{loc}$.

bessel: V^{PS} is simply derived from V_{eff} by a simple pseudization scheme using a zero-order spherical Bessel function: $V^{PS}(r) = \alpha \cdot \frac{\sin(q \cdot r)}{r}$ for $r < r_{vloc}$.

In that case, l_{loc} and E_{loc} are ignored and can be omitted.

Default (if missing) is **troulliermartins**

l_{loc} E_{loc} :

l quantum number and reference energy (Rydberg units) for use when

$Vloc_scheme = \mathbf{troulliermartins}$

or

$Vloc_scheme = \mathbf{ultrasoft}$

rc_1
 rc_2
...
 rc_{basis_size}

If Projector_keyword \neq "Blochl"

One line for each partial-wave

rc_i : (atomic units)

Matching radius used to get pseudo partial-waves $\tilde{\phi}_i(r)$ from partial-wave $\phi_i(r)$


As many radii as partial-waves have to be entered (one per line).

If projector_keyword is **blochl** or **VNCT**, these radii DO NOT HAVE TO BE GIVEN. In that case, they all are taken as $rc = r_{paw}$.

3. Test configurations

In the rest of the input file, the user can give electronic test configurations in order to test the validity of the created PAW dataset. For each electronic configuration the PAW Hamiltonian will be solved and resulting states printed.

Each test configuration has to be given as follow:



To end the list of configurations, enter a line with a 0 (zero):

0

Tests configurations are not mandatory and one can simply enter a "0" without having given any configuration.

Examples

A “minimal” input file

Boron [$1s^2$] $2s^2 2p^1$
4 partial-waves in basis

```
B 5.  
LDA-PW  
2 2 0 0 0  
2 1 1.0  
0 0 0  
c  
v  
v  
1  
1.7  
Y  
3.  
n  
Y  
3.  
n  
vanderbilt  
2 0.  
1.5  
1.5  
1.7  
1.7  
0
```

A "complete" input file

Nickel [$1s^2 2s^2 2s^6 3s^2 3p^6$] $3d^9 4s^1 4p^0$
6 partial-waves in basis

```
Nickel 28.
GGA-PBE scalarrelativistic point-nucleus loggrid 1500 80. 2.3
4 4 3 0 0 ! Up to 4s, 4p and 3d
1 0 2.0 ! Electronic configuration 3d9 4s1 4p0
2 0 2.0
2 1 6.0
3 0 2.0
3 1 6.0
3 2 9.0
4 0 1.0
4 1 0.0
0 0 0
c ! 1s
c ! 2s
c ! 3s
v ! 4s valence
c ! 2p
c ! 3p
v ! 4p valence
v ! 3d valence
2 ! Basis contains s, p and d partial-waves
2.3 2.3 1.1 2.2 ! rpaw=2.3, rshape=2.3, rveff=1.1, rcore=2.2
y ! Additional s partial-wave
4. ! at Eref=4.0 Ry
n
y ! Additional p partial-wave
4. ! at Eref=4.0 Ry
n
y ! Additional d partial-wave
2.5 ! at Eref=2.5 Ry
n
custom rrkj gramscmidtortho sinc ! RRKJ PW + sinc shape func.
Bessel ! Simple Bessel Vloc
2.3 ! Matching radius for Phi1 (l=0)
2.3 ! Matching radius for Phi2 (l=0)
2.3 ! Matching radius for Phi3 (l=1)
2.3 ! Matching radius for Phi4 (l=1)
2.3 ! Matching radius for Phi5 (l=2)
2.3 ! Matching radius for Phi6 (l=2)
1
1 0 2.0 ! Test configuration 3d8 4s2
2 0 2.0
2 1 6.0
3 0 2.0
3 1 6.0
3 2 8.0
4 0 2.0
0 0 0
0 ! END
```

Advice for use

In the following we give some keys for non-experienced users so that they can build input files for *atompaw* for new materials.

Short write-up

The first advice is to begin with a simple expression of the input file, setting most of the keywords to their default values.

- Concerning the all-electrons atomic computation, prefer a logarithmic grid ; test the influence of the number of grid points and, in case of difficulties, choose a regular grid. Begin with a scalar-relativistic solution of the wave equation. If the system shows convergence problems, try non-relativistic choice (not recommended when Z becomes high).
- Concerning the partial-waves basis generation, simply begin with:
 - a unique radius r_{paw}
 - 2 partial-waves per l angular momentum (if r_{paw} is small enough, 1 wave per l may suffice)
 - “bloechl” choice for `projector_keyword`
 - a norm-conserving *Troullier-Martins* pseudopotential at $l_{loc}=l_{max}+1$ and $E_{loc}=0$.

This choice should give a “stable” PAW dataset with correct physical results ; but Blöchl’s scheme for projectors can produce “inefficient” datasets (in the sense that they may need a large number of plane waves to converge the DFT calculation). To increase performance, choose the “vanderbilt” option for `projector_keyword`. The gain can be noticeable. But, generally, the best choice (for performance) would be “custom `rrkj`” projectors.

- Concerning the pseudopotential $V^{PS}(r)$, norm-conserving *Troullier-Martins* is generally the best choice but it can produce “ghost states” for d and f materials. If this happens, a simple “Bessel” pseudopotential can solve the problem. But, in the later case, one has to noticeably decrease the matching radius r_{vloc} (try $0.6*r_{paw}$ first).
- The other keywords in the input file can be adjusted (by experienced users) in order to obtain better “physical” results (by comparison with all-electrons calculations and/or experimental measurements)...

Detailed write-up

Here is a proposal for using *atompaw* to build new PAW datasets (from scratch). The procedure detailed here should help the user to generate optimal datasets in most cases.

☑ In a first stage, edit a simple input file for *atompaw*.

○ In the all-electrons atomic computation part

- ~ Define the material in the first line
- ~ Choose the exchange-correlation functional (LDA-PW or GGA-PBE) and select a scalar-relativistic wave equation and a (2000 points) logarithmic grid (second line).

“scalar-relativistic” is recommended for high Z materials

- ~ Then define the electronic configuration ; an excited configure may be useful if the PAW dataset is intended for use in a context where the material is charged (such as oxides).

Although, in our experience, the results are not highly dependent on the chosen electronic configuration.

- ~ Select the core and valence electrons: in a first approach, select only electrons from outer shells. But, if particular thermo dynamical conditions are to be simulated, it is generally needed to include “semi-core states” in the set of valence electrons. Semi-core states are generally needed with transition metal and rare-earth materials. There are also some cases (such as P) where physical conditions do not indicate a need for semi-core states, but the use of semi-core states are needed to avoid the appearance of the dreaded ghost states.

Note that all wave functions designated as valence electrons will be used in the partial-wave basis.

○ In the partial-waves basis generation part

Begin with a simple scheme. Select most of the keywords at their default values.

- ~ Enter only one matching radius (r_{paw}). Select it to be slightly less than half the inter-atomic distance in the solid (as a first choice).
- ~ Add additional partial-waves if needed: choose to have 2 partial-waves per angular momentum in the basis (this choice is not necessarily optimal but this is the most common one; if r_{paw} is small enough, 1 partial-wave per l may suffice). As a first guess, put all reference energies for additional partial-waves to 0 Rydberg.
- ~ Select a “bloechl” projector scheme and a norm-conserving *Troullier-Martins* pseudopotential at $l_{loc}=l_{max}+1$ and $E_{loc}=0$. “bloechl” will probably be changed later to make the PAW dataset more efficient.

○ In the test configuration part

- ~ Add one test configuration; a good idea is to test (at least) the electronic configuration used in the all-electrons atomic computation part.

☑ At this stage, run *atompaw* !

The generated PAW dataset is a first draft. Several parameters have to be adjusted, in order to get accurate results and efficient DFT calculations.

1. The sensitivity of results to some parameters has to be checked.

- The radial grid:
Try to select 700 points in the logarithmic grid and check if any noticeable difference in the results appears. If yes, adjust the size of the grid (else, keep 700 points). If the results are difficult to get converged, try a regular grid...
*For use with the pwpaw code, linear or logarithmic grids can be used.
For use with the abinit code, the logarithmic grid is preferred.*
- The relativistic approximation of the wave equation:
scalar-relativistic option should give better results than non-relativistic one, but it sometimes produces difficulties for the convergence of the atomic problem (either at the all-electrons resolution step or at the PAW Hamiltonian solution step). If convergence cannot be reached, try a nonrelativistic calculation (not recommended for high Z materials).
- A summary of the atomic all-electrons computation and the PAW dataset properties can be found in the [Atom_name](#) file (*Atom_name* is the first parameter of the input file). A look at the different values of *evale* (valence energy) is important. All-electron value has to be as close to others as possible. *evale* has to be insensitive to grids parameters.

2. Have a look at the partial-waves, PS partial-waves and projectors.

Plot the wfn.i files in a graphical tool of your choice. You should get 3 curves per file: $\varphi_i(r)$, $\tilde{\varphi}_i(r)$ and $\tilde{p}_i(r)$

- The $\tilde{\varphi}_i(r)$ should meet the $\varphi_i(r)$ near or after the last maximum (or minimum). If not, it is preferable to change the value of the matching radius rc_i .
- The $\tilde{\varphi}_i(r)$ and $\tilde{p}_i(r)$ should have the same order of magnitude. If not, you can try to get this in three ways:
 - Change the matching radius rc_i for this partial-wave; but this is not always possible... spheres cannot have a large overlap in the solid...
 - Change the pseudopotential scheme (see later).
 - If there are two (or more) partial waves for the considered l angular momentum, including additional partial waves (unbound states):
Decreasing the magnitude of projector is possible by displacing the references energies. Moving the energies away from each other generally reduce the magnitude of projectors, but a too big difference between energies can lead to wrong logarithmic derivatives (see following chapter).
- The two first values of *evale* (valence energy) in the [Atom_name](#) file have to be close. If not, choices for projectors and/or partial waves certainly are not judicious.
- Example of difficulty with $\tilde{p}_i(r)$: when the amplitude of projectors becomes too large, *atompaw* can produce an error with the following message:

```
No convergence in boundsep
Followed by
Best guess of eig, dele = xxxxx  yyyyy
```

This happens during the PAW Hamiltonian resolution (which cannot be achieved). One can bypass the difficulty by generating “softer” projectors as explained just above.

3. Have a look at the logarithmic derivatives.

They are printed in the `logderiv.l` files. Each `logderiv.l` file correspond to l quantum number and contains the logarithmic derivative of the l -state, $d(\log(\Psi_l(E)))/dr$, computed for exact atomic problem and with the PAW dataset.

- The 2 curves should be superimposed as much as possible. By construction, they are superimposed at the two energies corresponding to the two l partial-waves. If the superimposition is not good enough, the reference energy for the second l partial-wave should be changed.
- Generally a discontinuity in the logarithmic derivative curve appears at $0 <= E_0 <= 4$ Rydberg. A reasonable choice is to choose the 2 reference energies so that E_0 is in between (if possible, i.e. if one the 2 partial-waves correspond to an unbound state).
- Too close reference energies produce “hard” projector functions. But moving reference energies away from each other can damage accuracy of logarithmic derivatives.
- Another possible problem is the presence of a discontinuity in the PAW logarithmic derivative curve at an energy where the exact logarithmic derivative is continuous. This generally shows the presence of a “ghost state”.

First, try to change to value of reference energies; this sometimes can make the ghost state disappear.

If not, it can be useful to:

- Change the pseudopotential scheme. Norm-conserving pseudopotentials are sometimes so deep (attractive near $r=0$) that they produce ghost states. A first solution is to change the l quantum number used to generate the norm-conserving pseudopotential. But this is generally not sufficient. Changing the pseudopotential scheme is (in most cases) the only efficient cure.

Select a simple “`bessel`” pseudopotential can solve the problem. But, in that case, one has to noticeably decrease the matching radius r_{vloc} if one wants to keep reasonable physical results. Loosing too much norm for the wave function associated to the pseudopotential can have dramatic effects on the results.

Selecting a value of r_{vloc} between $0.6*r_{paw}$ and $0.8*r_{paw}$ is a good choice; but the best way to adjust r_{vloc} value is to have a look at the two first values of `evale` in `Atom_name` file. They have to be as equal as possible and are sensitive to the choice of r_{vloc} .

- Change the matching radius rc_i for one (or both) l partial-wave(s). In some cases, changing rc_i can remove ghost states ...

*In most cases (changing pseudopotential or matching radius), one has to **restart the procedure from step 2.** (only for l partial-waves).*

4. Now, one has to test the efficiency of the generated PAW dataset.

Run a DFT computation and determine the size of the plane wave basis needed to get a given accuracy. If the cut-off energy defining the plane waves basis is too high (higher than 20 Hartree, if matching radius has a reasonable value), some changes have to be made in the input file.

- First possibility: change `projector_keyword="bloechl"` by `projector_keyword="vanderbilt"`. Vanderbilt projectors generally are more localized in reciprocal space than Bloechl ones.
Recheck the plane waves cut-off (in a DFT calculation)... it should have decrease (but this is not a general rule).
- Second possibility: use RRKJ pseudization for PS partial-waves (put `projector_keyword="custom"` and `ps_keyword="rrkj"`). This pseudization is particularly efficient and gives highly localized projectors (in reciprocal space). This choice has, in most cases, the best influence on the plane wave basis.

One has to note that:

- ~ The localization of projectors in reciprocal space can (generally) be predicted by a look at `tprod.i` files. Such a file contains the curve of $q^2 \cdot \tilde{p}_i(q) \cdot \tilde{\varphi}_i(q)$ as a function of q (reciprocal space variable). q is given in Bohr⁻¹ units; it can be connected to the plane waves cut-off energy (in Hartree units) by: $E_{cut} = q_{cut}^2 / 2$. These quantities are only calculated for the bound states, since the Fourier transform of an extended function is not well-defined.
- ~ Generating projectors with Blöchl's scheme often gives the guaranty to have stable calculations. `atompaw` ends without any convergence problem and DFT calculations run without any divergence (but they need high plane wave cut-off). Vanderbilt projectors (and even more "custom" projectors) sometimes produce instabilities during the PAW dataset generation process and/or the DFT calculations...

In most cases, after having changed the projector generation scheme, one has to restart the procedure from step 2.

5. Finally, have a careful look at physical quantities obtained with the PAW dataset.

It can be useful to test their sensitivity to some input parameters:

- The analytical form and the cut-off radius r_{shape} of the shape function used in compensation charge density definition. By default a "sinc" function is used but "gaussian" shapes can have an influence on results. "Bessel" shapes are efficient and generally need a smaller cut-off radius ($\sim 0.8 \cdot r_{paw}$).
- The matching radius r_{core} used to get pseudo core density from atomic core density.
- The integration of additional ("semi-core") states in the set of valence electrons.
- The pseudization scheme used to get $V^{PS}(r)$.

All these parameters have to be meticulously checked, especially if the PAW dataset is used for non-standard solid structures or thermo dynamical domains.